

12

# EUROPEAN PATENT APPLICATION

21 Application number: 84301996.9

22 Date of filing: 23.03.84

51 Int. Cl.<sup>3</sup>: **C 12 N 15/00**  
 C 12 P 21/00, A 61 K 39/00  
 A 61 K 39/395, A 61 K 47/00  
 B 01 D 15/08, G 01 N 33/54  
 //C12R1/19, C12R1/125,  
 C12R1/865

30 Priority: 25.03.83 GB 8308235

43 Date of publication of application:  
 03.10.84 Bulletin 84/40

84 Designated Contracting States:  
 AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: **CELLTECH LIMITED**  
 244-250 Bath Road  
 Slough Berkshire SL1 4DY(GB)

72 Inventor: **Boss, Michael Alan**  
 20 Hunter Court Huntercombe Lane North  
 Slough Berkshire SL1 6DS(GB)

72 Inventor: **Kenten, John Henry**  
 24 Avery Avenue  
 Downley High Wycombe Buckinghamshire(GB)

72 Inventor: **Emtage, John Spencer**  
 12, Benjamin House Amersham Hill  
 High Wycombe Buckinghamshire(GB)

72 Inventor: **Wood, Clive Ross**  
 Whych Wood Buddle Hill  
 North Gorley NEAR Fordingbridge Hants(GB)

74 Representative: **Votier, Sidney David et al,**  
 CARPMAELS & RANSFORD 43, Bloomsbury Square  
 London WC1A 2RA(GB)

54 Multichain polypeptides or proteins and processes for their production.

57 This invention relates to multichain polypeptides or proteins and processes for their production in cells of host organisms which have been transformed by recombinant DNA techniques.

According to a first aspect of the present invention, there is provided a process for producing a heterologous multichain polypeptide or protein in a single host cell, which comprises transforming the host cell with DNA sequences coding for each of the polypeptide chains and expressing said polypeptide chains in said transformed host cell.

According to another aspect of the present invention there is provided as a product of recombinant DNA technology an Ig heavy or light chain or fragment thereof having an intact variable domain.

The invention also provides a process for increasing the level of protein expression in a transformed host cell and vectors and transformed host cells for use in the processes.




FIG. 1

MULTICHAIN POLYPEPTIDES OR PROTEINS AND PROCESSES FOR  
THEIR PRODUCTION

---

5 This invention relates to multichain polypeptides or  
proteins and processes for their production in cells  
of host organisms which have been transformed by recombi-  
nant DNA techniques.

10 In recent years advances in molecular biology based  
on recombinant DNA techniques have provided processes  
for the production of heterologous (foreign) polypeptides  
or proteins in host cells which have been transformed  
with heterologous DNA sequences which code for the  
production of these products.

15 Theoretically, the recombinant DNA approach may be  
applied to the production of any heterologous polypeptide  
or protein in a suitable host cell, provided that appro-  
priate DNA coding sequences can be identified and used  
20 to transform the host cell. In practice, when the  
recombinant DNA approach was first applied to the produc-  
tion of commercially useful products, its application  
for the production of any specified polypeptide or  
protein presented particular problems and difficulties,  
25 and the success of applying this approach to the production  
of any particular polypeptide or product was not readily  
predictable.

30 However, a large number of heterologous single chain  
polypeptides or proteins have now been produced by  
host cells transformed by recombinant DNA techniques.  
Examples of such heterologous single chain polypeptides  
or proteins include human interferons, the A and B  
chains of human insulin, human and bovine growth hormone,  
35 somatostatin, calf prochymosin and urokinase. Such

transformed host cells provide a reproducible supply of authentic heterologous polypeptide or protein which may be produced on an industrial scale using industrial fermentation technology.

5

It should be pointed out that some of these polypeptides, for instance urokinase, after secretion by a host cell appear as two chain molecules. However, in such cases, the molecule is synthesised by the host cell as a single chain polypeptide, coded for by a single DNA sequence, which is cleaved in the host cell subsequent to synthesis to form the two chain structure.

It is known that in both human and animal systems there are a number of polypeptides or proteins which have multichain structure in which the chains are not derived from the cleavage of a single chain polypeptide coded for by a single DNA sequence. In such cases, the gene for each of the chains may be located at different points on the same chromosome or even on different chromosomes. In these cases, the polypeptide chains are synthesised separately and then assembled into the complete molecule subsequent to synthesis. Heretofore, no such multichain polypeptide or protein has been produced by recombinant DNA techniques from a single host cell.

A particular example of a class of such multichain polypeptides or proteins is the immunoglobulins.

Immunoglobulins, commonly referred to as antibodies, are protein molecules produced in animals by B-lymphocyte cells in response to challenge with foreign antigenic agents, such as bacteria, viruses and foreign proteins. The immunoglobulins comprise a crucial part of the immune systems of humans and animals. The immunoglobulins

recognise specific parts of the foreign agents and bind onto them. The specific parts are usually known as antigenic determinants or antibody binding sites. A given foreign agent is likely to have a number of different antigenic determinants.

A typical immunoglobulin (Ig) molecule is shown in Figure 1 of the accompanying drawings, to which reference is now made. The Ig molecule comprises two identical polypeptide chains of about 600 amino acid residues (usually referred to as the heavy chains H), disulphide bonded to each other, and two identical shorter polypeptide chains of about 220 amino acid residues (usually referred to as the light chains L), each light chain being disulphide bonded to one end of each heavy chain as shown.

When the Ig molecule is correctly folded, each chain is formed into a number of distinct globular areas, usually known as domains, joined by a more linear polypeptide chain. The light chains have two such domains, one of which is of variable sequence  $V_L$ , and the other of which is of constant sequence  $C_L$ . The heavy chains have a single variable domain  $V_H$  adjacent the variable domain  $V_L$  of the light chain, and three or four constant domains  $C_{H1-3}$  or 4.

The variable domains on both the heavy and the light chains each contain three hypervariable regions (HV1-3) which, when the Ig molecule is correctly folded, are located adjacent one another and form the antigen binding site. It is this area which recognises and binds to the antigenic determinant for which the Ig molecule is specific.

The constant domains of the Ig molecule do not take part in the binding to the antigenic determinant, but mediate the actions triggered by the binding of the Ig molecule to the antigenic determinant. It is believed that this triggering is caused by an allosteric effect induced by the binding of the Ig molecule to the antigenic determinant. The constant domain may enable the Ig molecule to fix complement or may cause mast cells to release histamine.

Ig's may be categorised by class or subclass, depending on which of a number of possible heavy chain constant domains they contain, there being eight such possible heavy chains in mice. Thus, for instance, an Ig molecule with a  $\mu$  heavy chain belongs to the class IgM, and one with a  $\gamma_1$  heavy chain to the class IgG<sub>1</sub>.

Ig's may also contain one of two light chains, designated as  $\kappa$  and  $\lambda$  light chains, which have different constant domains and different sets of variable domains.

The structure of the Ig molecule and the location of the genes coding for the various domains thereof are discussed more fully by Early and Hood, in Genetic Engineering, Principles and Methods, Vol.3, pages 153-158 (edited by Setlow and Hollaender, Plenum Press).

It is known that Ig molecules on digestion with selected enzymes can produce a number of immunologically functional fragments. Two such fragments are known as the  $F_{ab}$  and  $(F'_{ab})_2$  fragments. The  $F_{ab}$  fragment comprises one light chain linked to the  $V_H$  and  $C_{H1}$  domains of a heavy chain as shown in Figure 1. The  $(F'_{ab})_2$  fragment consists essentially of two  $F_{ab}$  fragments linked together by a small additional portion of the heavy chains as shown

in Figure 1. These fragments and other similar fragments can be of use in various tests and diagnostic and medical methods.

5 The principle method employed for the production of Ig's involves the immunisation of susceptible animals with the antigenic agents to provide an immune reaction. Generally the animal is immunised a second time to improve the yield of Ig. The animal is then bled and  
10 the Ig is recovered from the serum.

However, the product of this method is not a homogeneous protein. The animal will produce Ig of different classes and also Ig specific for each of the different antigenic  
15 determinants on the antigenic agent, and its blood will therefore contain a heterogeneous mixture of Ig's. Obtaining a specific Ig of particular class and desired specificity from such a mixture requires very difficult and tedious purification procedures.

20

Recently, it has become possible to produce a homogeneous Ig of a single class and a single specificity by a technique first described by Kohler and Milstein (Nature, 256, 495-479, 1975). The technique involves the fusion  
25 of single Ig-producing parent cells with cancer cells to produce a monoclonal hybridoma cell which produces the Ig. Ig produced by this technique is usually known as monoclonal antibody. The nature of the monoclonal Ig is determined by the class and specificity of the Ig  
30 produced by the parent cell.

Recently, attempts have been made to use recombinant DNA techniques to produce fragments of Ig molecules. For instance, Amster et al. (Nucleic Acid Research, 8,  
35 No.9, 1980, pp 2055 to 2065) disclose the cloning of

doubl stranded cDNA sequences encoding for a mouse Ig light chain into a plasmid. An E. Coli strain transformed by the plasmid synthesis d a protein thought to comprise the complete constant domain of the light chain and  
5 about 40 amino acid residues of its variable region.

Kemp and Cowman (Proc. Natl. Acad. Sci. USA, 78, 1981, pp 4520 to 4524) disclose the cloning of cDNA sequences encoding for mouse heavy chain fragments and the trans-  
10 forming of an E. Coli strain which then synthesised heavy chain polypeptide fragments.

In both these cases, the polypeptides were produced as fusion proteins, in which the fragments of the Ig poly-  
15 peptides were fused with additional non-Ig polypeptide sequences, and with incomplete variable domains. Thus, the polypeptide chains produced in these studies were not immunologically functional polypeptides as they were incapable of combining with complementary heavy  
20 or light chains to provide Ig molecules having intact antigen binding sites and immunological function.

Research studies have also been carried out in mammalian systems. For instance, Falkner and Zachau (Nature, 298,  
25 1982, pp 286 to 288) report the cloning of cDNA sequences encoding for mouse light chains into a plasmid which was used to transfect genomic eukaryotic cells which could then transiently synthesise light chains.

30 Rice and Baltimore (Proc. Natl. Acad. Sci. USA, 79, 1982, pp 7862 to 7865) report on the transfection of a functionally rearranged K light chain Ig gene into a murine leukemia virus-transformed lymphoid cell line. The cell line is then able to express the gene continuously.

35 In both these cases, the K genes used to transfect the

mammalian cells were obtained from myeloma cells and the K polypeptides produced were of indeterminate immunological function.

- 5 A further approach is exemplified in a series of papers by Valle et al. (Nature, 291, 1981, 338-340; Nature, 300, 1982, 71-74; and J. Mol. Biol., 160, 1982, 459-474), which describe the microinjection of mRNAs encoding for heavy or light chains of Ig isolated from a mouse myeloma
- 10 line into oocytes of *Xenopus laevis*. Under certain conditions complete Ig molecules were formed. However, the mRNAs were obtained from myeloma cells and the Ig molecules were of indeterminate immunological function.
- 15 It can thus be seen that hitherto it has not been possible to produce functional Ig by recombinant DNA technology.

According to a first aspect of the present invention, there is provided a process for producing a heterologous

20 multichain polypeptide or protein in a single host cell, which comprises transforming the host cell with DNA sequences coding for each of the polypeptide chains and expressing said polypeptide chains in said transformed host cell.

25 According to a second aspect of the present invention, there is provided a heterologous multichain polypeptide or protein produced by recombinant DNA technology from a single host cell.

30 The present invention is of particular, but not exclusive, application in the production of Ig molecules and immunologically functional

---



Ig fragments of the type referred to above. However, it will be appreciated that the present invention can be applied to the production of other multichain polypeptides or proteins.

5

In relation to the product of Ig molecules according to the invention it will be appreciated that, in order to produce a functional molecule, the DNA sequences used to transform the host cell will need to encode  
10 for at least the  $V_L$  and  $V_H$  domains of an Ig molecule. Moreover, these domains will need to be complementary so that when the two polypeptide chains fold together they form an antigen binding site of predetermined specificity.

15

Preferably, the Ig molecule or fragment includes a complete light chain and at least the  $C_{H1}$  domain in addition to the  $V_H$  domain of the heavy chain. Most preferably the Ig molecule is intact.

20

It has also been shown by the present applicants that it is now possible to produce individual heavy and light chains having intact variable domains. This has not previously been possible. Therefore,  
25 according to a third aspect of the present invention there is provided as a product of recombinant DNA technology an Ig heavy or light chain or fragment thereof having an intact variable domain.

30 Advantageously, the Ig molecule or functional fragment thereof according to the present invention has a variable region (formed by the  $V_L$  and  $V_H$  domains) which defines a binding site for an antigenic determinant of clinical or industrial importance. The DNA  
35 coding sequences necessary to produce such a molecule may be derived from naturally occurring or hybridoma (monoclonal) Ig-producing cells with the desired specificity.

The constant domains of the Ig molecule or fragment, if present, may be derived from the same cell line as the variable region. However, the constant domains may be specifically altered, partially or completely omitted, or derived from a cell line producing a different class of Ig to provide Ig molecules or fragments having desired properties.

For example, an Ig molecule may be produced having variable domains ( $V_H$  and  $V_L$ ) identical with those from a monoclonal antibody having a desired specificity, and constant domain(s) from a different monoclonal antibody having desired properties, for instance to provide human compatibility or to provide a complement binding site.

Such alterations in the amino acid sequence of the constant domains may be achieved by suitable mutation or partial synthesis and replacement or partial or complete substitution of appropriate regions of the corresponding DNA coding sequences. Substitute constant domain portions may be obtained from compatible recombinant DNA sequences.

The invention may be utilised for the production of Ig molecules or fragments useful for immunopurification, immunoassays, cytochemical labelling and targetting methods, and methods of diagnosis or therapy. For example, the Ig molecule or fragment may bind to a therapeutically active protein such as interferon or a blood clotting factor, for example Factor VIII, and may therefore be used to produce an affinity chromatography medium for use in the immunopurification or assay of the protein.

It is also envisaged that the Ig molecule may be synthesised by a host cell with another peptide moiety attached to one of its constant domains. Such a further peptide moiety may be cytotoxic or enzymatic. Alternatively, the moiety may be useful in attaching the Ig molecule to a biological substrate, such as a cell or tissue, or to a non-biological substrate, such as a chromatography medium. Such a peptide moiety is herein referred to as a structural peptide moiety.

It is further envisaged that cytotoxic, enzymic or structural peptide moieties could be attached to the Ig molecule by normal peptide chemical methods, as are already known in the art, rather than by being synthesised with the Ig molecule.

The Ig molecule or fragment may also comprise a therapeutic agent in its own right. For instance, an Ig molecule or fragment specific for D blood group antigen may be useful for the prevention of haemolytic disease of the new born.

Any suitable recombinant DNA technique may be used in the production of the multichain polypeptides or proteins of the present invention. Typical expression vectors such as plasmids are constructed comprising DNA sequences coding for each of the chains of the polypeptide or protein.

It will be appreciated that a single vector may be constructed which contains the DNA sequences coding for more than one of the chains. For instance, the DNA sequences coding for Ig heavy and light chains may be inserted at different positions on the same plasmid.

Alternatively, the DNA sequence coding for each chain may be inserted individually into a plasmid, thus producing a number of constructed plasmids, each coding for a particular chain. Preferably the plasmids into which the sequences are inserted are compatible.

The or each constructed plasmid is used to transform a host cell so that each host cell contains DNA sequences coding for each of the chains in the polypeptide or protein.

Suitable expression vectors which may be used for cloning in bacterial systems include plasmids, such as Col E1, pCR1, pBR322, pACYC 184 and RP4, phage DNA or derivatives of any of these.

For use in cloning in yeast systems, suitable expression vectors include plasmids based on a 2 micron origin.

Any plasmid containing an appropriate mammalian gene promoter sequence may be used for cloning in mammalian systems. Such vectors include plasmids derived from, for instance, pBR322, bovine papilloma virus, retroviruses, DNA viruses and vaccinia viruses.

Suitable host cells which may be used for expression of the heterologous multichain polypeptide or protein include bacteria, such as E. Coli and B. Subtilis, streptomyces, yeasts, such as S. cervisiae, and eukaryotic cells, such as insect or mammalian cell lines. Examples of suitable bacterial host cells include E. Coli HB 101, E. Coli X1776, E. Coli X2882, E. Coli PS 410, E. Coli MRC 1, E. Coli RV308, E. Coli E103S and E. Coli B.

The present invention also includes constructed expression vectors and transformed host cells for use in producing the multichain polypeptides or proteins of the present invention.

5

After expression of the individual chains in the same host cell, they may be recovered to provide the complete multichain polypeptide or protein in active form, for instance to provide an Ig molecule of predetermined immunological function.

10

It is envisaged that in preferred forms of the invention, the individual chains will be processed by the host cell to form the complete polypeptide or protein which advantageously is secreted therefrom.

15

However, it may be that the individual chains may be produced in insoluble or membrane-bound form. It may therefore be necessary to solubilise the individual chains and allow the chains to refold in solution to form the active multichain polypeptide or protein. A suitable procedure for solubilising polypeptide chains expressed in insoluble or membrane-bound form is disclosed in our copending application No. (Protein Recovery, Agent's Ref. GF 402120 and 402121).

20

It will be appreciated that the present application shows for the first time that it is possible to transform a host cell so that it can express two or more separate polypeptides which may be assembled to form a complete multichain polypeptide or protein. There is no disclosure or suggestion of the present invention in the prior art, which relates solely to the production of a single chain heterologous polypeptide or protein from each host cell.

30

35

The present invention will now be described, by way of example only, with reference to the accompanying drawings, in which:-

- Figure 1 shows a diagrammatic representation of a typical  
5 intact Ig molecule;  
Figure 2 shows the construction of plasmids for the direct synthesis of a  $\lambda$  light chain in E. Coli;  
Figure 3 shows the construction of plasmids for the direct synthesis of a  $\mu$  heavy chain in E. Coli;  
10 Figure 4 is a diagrammatic representation of  $\mu$  mRNA sequences around the initiation codon;  
Figure 5 shows the construction of plasmids having altered secondary structure around the initiation codon;  
15 Figure 6 is a polyacrylamide gel showing expression and distribution of  $\mu$  protein from E. Coli B;  
Figure 7 is a polyacrylamide gel showing pulse chase autoradiograms of  $\mu$  protein in E. Coli B and in E. Coli HB101;  
20 Figure 8 is a polyacrylamide gel showing the results of  $\lambda$  gene expression in E. Coli;  
Figure 9 is a polyacrylamide gel showing the distribution of recombinant  $\lambda$  light chain polypeptide between the soluble and insoluble cell fractions;  
25 Figure 10 is a polyacrylamide gel showing expression and distribution of  $\lambda$  protein from E. Coli El03S;  
Figure 11 shows the results of the fractionation of  $\mu$  and  $\lambda$  protein expressed by E. Coli B on DEAE Sephacel;  
Figure 12 shows the specific hapten binding of reconstituted Ig molecules; and  
30 Figure 13 shows the heteroclitic nature of the hapten binding of the reconstituted Ig molecules.

In the following examples, there is described the production of Ig light and heavy chain polypeptides derived from monoclonal antibodies which recognise and bind to the antigenic determinant 4-hydroxy-3-nitrophenyl acetyl (NP), using E.coli and S. cerevisiae as the host cells. Recombinant DNA techniques were used to enable the host cells to express both the polypeptide chains.

It will be appreciated that the invention is not limited to the specific methods and construction described hereafter.

#### Construction of Lambda Light Chain Expression Plasmid

Figure 2, to which reference is now made, shows schematically the method used to construct a  $\lambda$  light chain expression plasmid.

It was decided to express the lambda gene in E. coli by direct expression of the gene lacking the eucaryotic signal peptide but containing a methionine initiator residue at the amino-terminus (met-lambda). The approach used for bacterial synthesis of met-lambda was to reconstruct the gene in vitro from restriction fragments of a cDNA clone and to utilise synthetic DNA fragments for insertion into the bacterial plasmid pCT54 (Emtage et al., Proc. Natl. Acad. Sci. USA., 80, 3671 to 3675, 1983). This vector contains the E. coli trp promoter, operator and leader ribosome binding site; in addition 14 nucleotides downstream of the ribosome binding site is an initiator ATG followed immediately by EcoRI and HindIII sites and the terminator for E. coli RNA polymerase from bacteriophage T7.

As a source of light chain we used a plasmid pAB $\lambda$ 1-15 which contains a full-length  $\lambda_1$  light chain cDNA cloned into the PstI site of pBR322. This  $\lambda_1$  light chain is derived from a monoclonal antibody, designated S43, which binds to 4-hydroxy-3-nitrophenylacetyl (NP) haptens.

In order to create a HindIII site 3' to the end of the lambda gene for insertion into the HindIII site of pCT54, the cDNA was excised from pAB $\lambda$ 1-15 using PstI. The cohesive ends were blunt ended using the Klenow fragment of DNA polymerase and synthetic HindIII linker molecules of sequence 5'-CCAAGCTTGG-3' ligated. The DNA was digested with HindIII and the 850bp lambda gene isolated by gel electrophoresis and cloned into HindIII cut pAT153 to yield plasmid pAT $\lambda$ 1-15. The 3' end of the lambda gene was isolated from pAT $\lambda$ 1-15 by HindIII plus partial SacI digestion as a 630bp SacI-HindIII fragment (2 in Figure 2). The HindIII cohesive end was dephosphorylated by calf intestinal alkaline phosphatase during isolation of the fragment to prevent unwanted ligations at this end in subsequent reactions.

A HinfI restriction site is located between codons 7 and 8 and the lambda sequence. The 5' end of the lambda gene was isolated as a 148bp HinfI to SacI fragment (1 in Figure 2).

Two oligodeoxyribonucleotides were designed to restore codons 1-8, and to provide an initiator ATG as well as BclI and HinfI sticky ends. The two chemically synthesised oligonucleotides made to facilitate assembly of the gene had the sequences:

R45 5' -pGATCAATGCAGGCTGTTGTG 3'

R44 3' CCGACAACACTGAGTCCTTAp- 5'

pCT54 was cut with both BclI and HindIII and the resulting linear molecules isolated, mixed together with the two oligodeoxyribonucleotide linkers R44 and R45 and both fragments 1 and 2, and ligated using T4 ligase (Figure 2). The mixture was used to transform E. coli DH1 to ampicillin resistance. Recombinant clones in pCT54 were identified by hybridisation of DNA from replica plated colonies on nitrocellulose to a nick-translated probe derived from the pAT $\lambda$ 1-15 insert.




A clone was identified which hybridised to lambda cDNA and also showed the predicted restriction fragment pattern. This plasmid (designated pCT54 19-1) was sequenced from the ClaI site and shown to have the anticipated sequence

- 5 except that there was a mutation of the fourth codon from CTG to ATG, changing the amino acid at this point from valine to methionine.

The sequence in this area was:

...GATTGATCA.ATG.CAG.GCT.GTT.ATG.ACT.CAG.GAA.TCT.GCA.CTC.ACC.ACA.TCA..

met gln ala val met thr gln glu ser ala leu thr thr ser

- 10 The restriction enzyme sites in pCT54 between the Shine and Dalgarno sequence (AAGG), which is important for ribosome binding, and the ATG allow for the adjustment of the SD-ATG distance, an important parameter in determining expression rates. The SD-ATG distance was reduced by
- 15 cutting the plasmid with ClaI or BclI and creating blunt ended species by digestion with  $S_1$  nuclease. 2  $\mu$ g of ClaI cut DNA was digested with 200 units of  $S_1$  nuclease for 30 minutes at 30° using standard buffer conditions. The solution was deproteinised with phenol and the DNA re-
- 20 covered by ethanol precipitation. This DNA on religation with T4 DNA ligase and transformation into E.coli strain HB101 gave rise to a number of plasmids which had lost the ClaI or BclI site.
- 

The plasmids which had lost their ClaI site were sequenced in the region surrounding the initiator ATG.

....AAGGGTATTGATCAATG CAG.... plasmid pNP3

SD met glu

....AAGGGTTTGATCAATG CAG plasmid pNP4

SD met glu

In order to achieve high level expression a number of other approaches were followed. Firstly, a series of constructs were obtained which had increasing amounts of the 3' untranslated region of the cDNA removed by Bal 31 exonuclease. Secondly, a high copy number plasmid containing  $\lambda$  cDNA was constructed. This plasmid contained a par function (Meacock, P. A. and Cohen, S. N. Cell, 20, 529-542, 1980) as well as being present in high copy number. Thirdly, the pNP3 plasmid was transformed into a number of protease-deficient strains or into HB101 in conjunction with a protease deficient dominant acting plasmid (Grossman, A. D. et al Cell, 32, 151-159, 1983).

#### Construction of $\mu$ Heavy Chain Expression Plasmid


The full-length  $\mu$  heavy chain cDNA derived from the NP binding monoclonal antibody B1-8 had been cloned into the PstI site of pBR322 yielding a plasmid designated pAB $\mu$ -11 (Bothwell et al, Cell, 24, 625-637, 1981). In order to achieve high level expression, the  $\mu$  cDNA minus the eukaryotic leader was reconstructed into pCT54. The construction of the  $\mu$  heavy chain expression plasmid is shown diagrammatically in Figure 3. Two chemically synthesised oligonucleotides were made to facilitate this. These have the following sequences:

R43 5' GATCAATGCAGGTTTCAGCTGCA 3'

30 R46 3' TTACGTCCAAGTCG 5'

These were ligated into BclI cut pCT54 using T4 DNA ligase and the resulting plasmid designated pCT54 Pst. The linkers were designed to replace the sequence in pCT54 between the BclI site and the ATG, to provide an internal 5 PstI site and to recreate the sequence from pABu-11 5' to the PstI site up to codon +1. pCT54 Pst was cut briefly with PstI, treated with alkaline phosphatase and full length DNA glass isolated from a 1% agarose gel. Similarly pABu-11 was briefly cut with PstI and the full 10 length  $\mu$  insert isolated following agarose gel electrophoresis. The  $\mu$  cDNA was ligated with the full length pCT54Pst fragment using T4 DNA ligase under standard conditions and a plasmid designated pCT54 $\mu$  was identified which by restriction enzyme analysis was shown to contain 15 a full-length  $\mu$  insert. The plasmid was sequenced around the 5' linker region and was found to have the anticipated sequence:

5' ....TGATCAATGCAGGTTTCAGCTGCAGGGGGGGGATCGGATCGAG.... 3', demonstrating that it was, indeed, a full length clone. A 20 complete PstI digest of pCT54 $\mu$  liberated a 1.4 Kb fragment which was purified by 0.8% agarose gel electrophoresis and glass powder isolation. This was ligated with PstI cut pCT54 Pst (see above) using T4 DNA ligase under standard conditions, followed by transformation into HB101. A 25 plasmid designated pNP1 was isolated which was shown by restriction endonuclease pattern analysis to contain the 1.4 Kb  $\mu$ cDNA fragment in an appropriate orientation.



(Figure 3). pNP1 is a plasmid which consisted of an appropriate 5' end for expression, whilst pCT54 contained an appropriate 3' end. The full length gene was re-constructed into pCT54 by cutting both pNP1 and pCT54 with *Ava*I which cuts once in the plasmid and once in the  $\mu$  gene. Both digests were run on a 1% agarose gel and the 1.9 Kb fragment from pNP1 and the 3.65 Kb fragment from pCT54 isolated. Following alkaline phosphatase treatment of the 3.65 Kb pCT54 fragment, the two pieces of DNA were ligated to each other and transformed into HB101. A plasmid designated pNP2 was identified which demonstrated the correct restriction endonuclease pattern. It was sequenced in the area surrounding the initiator ATG and found to have the anticipated sequence:

5' ....TTGATCAATGCAGGTTTCAGCTGCAGCAGCCTGGGGCTGAGCTTGTGAAG.... 3'

The vector pCT54 had been constructed to include two restriction sites (*Bcl*I and *Cla*I) between the S-D sequence (AAGG) and the initiation codon so that the distance between these sequence elements could be varied. As most *E.coli* mRNAs have 6-11 nucleotides between the S-D sequence and the AUG, the distance in pNP2 was reduced by modification at the *Cla*I site. pNP2 was cut with *Cla*I and incubated with S1. The amount of S1 nuclease was adjusted so that some DNA molecules would lose 1-2 extra base pairs as a result of 'nibbling' by the enzyme. This DNA on religation with T4 DNA ligase and transformation into *E.coli* strain HB101 gave rise to a number of colonies harbouring plasmids which had lost the *Cla*I site. The sequences around the initiation codon of four plasmids pNP223, pNP261, pNP9 and pNP282 were determined and are given in Table 1, part A below.

TABLE 1

		S-D to ATG distance (no. of residues)	Sequence (5' - 3')	Expression (mu) (ru)		
				A	B	C
(A)	DNP2	14	-AAAAAGGSTATCGATTGATCAATGCAAGTTTCAG-	0.7	0.6	0.8
S1 derivatives of DNP2	DNP223	12	-AAAAAGGSTATATTGATCAATGCAAGTTTCAG-	-	0.7	0.6
	DNP261	10	-AAAAAGGSTATGATCAATGCAAGTTTCAG-	-	1.1	1.1
	DNP282	7	-AAAAAGGSGATCAATGCAAGTTTCAG-	-	0.7	0.6
	DNP9	9	-AAAAAGGSTATGATCAATGCAAGTTTCAG-	1.0	1.0	1.0
(B)	DNP11	11	-AAAAAGGSTATGACATATGCAAGTGCAG-	5.3	6.3	9.1
oligo constructs	DNP12	11	-AAAAAGGSTATGACATATGCAAGTTTCAG-	1.8	2.3	1.3
	DNP14	9	-AAAAAGGSTATGATCAATGCAAGTGCAG-	72.2	106.7	124.6
	DNP8	-	-	77.0	53.4	145.5
	DCT70	-	-	0	0	0

The expression of these constructs in *E. coli* B was examined in three experiments, and concentrations of  $\mu$  protein were determined by ELISA as relative units (ru), a measure of  $\mu$  concentration normalised to pNP9. The results are shown in Table 1.

Inductions were carried out by resuspending cells at 1:50 dilution from overnight cultures into induction medium, consisting of:  $\text{KH}_2\text{PO}_4$  (3g/L),  $\text{Na}_2\text{HPO}_4$  (6g/L), NaCl (0.5g/L), Difco vitamin assay casamino acids (30g/L),  $\text{NH}_4\text{Cl}$  (4g/L) glycerol (16g/L), proline (0.5g/L), Difco yeast extract (1g/L),  $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$  (0.022g/L),  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  (0.025g/L), thiamine (0.01g/L) and carbenicillin (0.1g/L). Cultures were shaken at 37°C and cells harvested by centrifugation.

ELISA assays were performed essentially as described by Patel et al. (Nucleic Acid Research, 10, 5605 to 5620, 1982), using affinity purified goat anti-mouse IgM (Tago) and affinity purified goat anti-mouse IgM (Tago) conjugated to peroxidase, with 3, 3', 5, 5' tetramethylbenzidine (Miles) as a substrate. Bacterial cell pellets were boiled for 2 minutes in 9M urea (BRL) and loaded directly into the ELISA, such that all wells contained 2.25M urea, 10mM Tris-HCl pH7.8, 150mM NaCl, 0.05% NP40 and 0.5% casein (Hammarsten, B.D.H.).

Altering the S-D to ATG distance with nuclease S1 was found to increase the level of  $\mu$  expression, relative to the parental plasmid pNP2, but only to a small extent (compare pNP9). The optimal S-D to AUG distance was found to be 9-10 nucleotides, as found in pNP9 and pNP261 respectively.

Secondary Structure Analysis

Hairpin loops were identified by use of the computer programme HAIRGU developed by Dr. Roger Staden. The  $\Delta G$  values were calculated as described by Trijico et al. (Nature NB, 246, 40-41, 1973).

Analysis of the potential secondary structure of  $\mu$  mRNA encoded by pNP2 and pNP9 (one of the S1 derivatives of pNP2) revealed an extensive array of possible hairpin loops. Attention was focused on the region of this mRNA containing the ribosome binding site since the sequestering of S-D sequences and initiation codons into secondary structure may inhibit the initiation of translation. It was found that a hairpin loop including the U and G of the initiation codon could be formed (see Figure 4). This hairpin loop is formed entirely within coding sequences and has a  $\Delta G = -7.6$  kcal. No secondary structures were found which buried the S-D sequence, and none which mutually excluded the hairpin loop described above, and which had a lower  $\Delta G$ .

To test whether expression of  $\mu$  protein could be influenced by changes designed to alter potential secondary structure in the ribosome binding region we used synthetic oligonucleotides to mutagenise this region of  $\mu$  mRNA. The plasmids were constructed by insertion of a pair of oligonucleotides between the blunted Cla I site and the 5' Pst site of  $\mu$  in pNP2 (see Figure 5). The cloning strategy used to accomplish this, however, was multi-staged due to the presence of four Pst sites in pNP2. The vector pACYC184CM, which lacks Cla I and Pst I sites, was first constructed by cutting pACYC184 at its unique Cla I site, 'filling-in' the cohesive termini using T4 DNA polymerase and religating. Next, pACYC184CM was cut with BamHI, and the BamHI fragment from pNP2, bearing the 5'  $\mu$  sequence, was inserted into it to form pCM $\mu$  (see Figure 4). The vector pCM $\mu$  contained unique Cla I and Pst sites which were required for the oligonucleotide cloning (see Figure 5).

pCM $\mu$  was cut with Cla I to completion, blunted with nuclease S1, and then cut with Pst I. The cut plasmid was purified by agarose gel electrophoresis and ligated separately with pairs of oligonucleotides, with the latter in a 100-fold molar excess. The pairs of oligonucleotides used were: R131 and R132 (for pNP11); R196 and R197 (for pNP12); and R202 and R203 (for pNP14). These have the sequences given below.

TGCACATATGCAAGTGCAACTGCA (R131), GTTGCACCTTGCATATGTGCA (R132),  
 GCTGAACCTGCATATGTGCA (R196), TGCACATATGCAGGTTTCAGCTGCA (R197),  
 GTTGCACCTTGCAATTGATC (R202), GATCAATGCAAGTGCAACTGCA (R203).

The ligation mixture was then used to transform HB101.

Recombinant clones were detected by colony hybridisation, on nitrocellulose filters, using one of each pair of oligonucleotides as a probe. Positive clones were sequenced and shown to have the correct nucleotide sequence. The  $\mu$  sequences containing the cloned oligonucleotides were excised on Eco RV-Bgl II fragments. These were ligated to Eco RV-Bgl II cut pNP2 to reconstruct the full length  $\mu$  genes (see Figure 5). Three different plasmids were created by these procedures; pNP11, pNP12 and pNP14, having the nucleotide sequences shown in Table 1 above.

In the construct pNP11 the potentially deleterious hairpin loop of pNP2 was abolished by changing the residues in the degenerate position of three codons and by introducing a different A-D to A-U sequence. These changes allow the region between the S-D and AUG to base pair with the 5' end of  $\mu$  coding sequence to form a hairpin loop of  $\Delta G = -7.8$  kcal, approximately equal to that of pNP2, but leaving the AUG and S-D sequences exposed (see Figure 4). The only other mutually exclusive hairpin loops are of  $\Delta G = -2.3$ ,  $-3.6$  and  $+0.4$  kcal.

The construct pNP12 retains the S-D to AUG sequence of pNP11, but has the coding sequence of pNP2, such that the hairpin loop of pNP2 can form (see Figure 4).

The final  $\mu$  construct was pNP14, which had the S-D to AUG sequence of pNP9, but the 5' coding sequence of pNP11, so that the hairpin loop of pNP12 could not form. The construct pNP14 had no secondary structure burying the S-D or  $\mu$  AUG.

#### $\mu$ Protein Expression from Constructs with Differing mRNA Secondary Structures

E.coli B strains containing plasmids pNP9, pNP11, pNP12 and pNP14 were grown in induction medium and samples removed for ELISA assay. The concentration of  $\mu$  was increased 6-7 fold in E.coli B cells containing pNP11 relative to pNP9 (see Table 1, Part b). Plasmid pNP12, which has the S-D to AUG sequence of pNP11, but the coding sequence of pNP9, showed only a two-fold increase in level of  $\mu$  expression compared to pNP9.



Plasmid pNP14, which has the S-D to AUG sequence of pNP9 but the 5'  $\mu$  coding sequence of pNP11, and thus differed from pNP9 only in three residues, was found to express  $\mu$  protein over 90 times that found in pNP9.

A plasmid, pNP8, has also been constructed, from which  $\mu$  is expressed,

5 also from the trp promoter, as part of a trpE- $\mu$  fusion protein containing the amino terminal 53 amino acids of trpE and the carboxyl terminal 503 amino acids of  $\mu$ . The rationale for making this fusion gene was that  $\mu$  would be translated from an efficient bacterial ribosome binding site rather than from one of unknown efficiency. Then, by comparing the amounts  
10 of trpE- $\mu$  produced from pNP8 with those of  $\mu$  produced from the different constructs, an estimate of relative ribosome binding efficiency may be made. When the trpE- $\mu$  gene in pNP8 was induced and the products quantitated by ELISA, it was found that the  $\mu$  expression in strains containing pNP8 and pNP14 were very similar (see Table 1, part B).

15 The ribosome binding site (RBS) sequences of pNP11, pNP12 and pNP14 were designed in order to test the hypothesis that sequestering of the initiation codon into secondary structure inhibits translation initiation, and to increase the rate of initiation by releasing the initiation codon. In pNP11, the RBS hairpin loop of pNP2 and pNP9 was abolished by changing the 5'  $\mu$  coding  
20 sequence in the degenerate position of three codons. In addition, the S-D to AUG sequence was changed to allow the base-pairing of this region with the 5' coding sequence. The result of these changes is that the initiation codon is exposed in the loop of a hairpin with the S-D sequence at the base of the stem (see Figure 4). This hairpin loop had a  
25  $\Delta G$  approximately equivalent to the  $\Delta G$  of the pNP9 hairpin loop. Strains containing pNP11 produced 6-7 times as much  $\mu$  as did those with pNP9. To test whether this was due to the change in S-D to ATG sequence characteristics, rather than to the liberation of the initiation codon, pNP12 was constructed as a control. The latter has the S-D to AUG sequence  
30 of pNP11, but retains the 5'  $\mu$  coding sequence of pNP9, so that the RBS hairpin loop of pNP11 cannot form, but the potentially deleterious one of pNP9 can form. The level of  $\mu$  expression from pNP12 was found to be increased by only two-fold relative to pNP9. This increase is probably due to an altered S-D to AUG distance, a U immediately 5' to the  
35 initiation codon , and the S-D to AUG sequence being G-poor each of which has been shown to be advantageous. It is therefore likely that

the increased expression from pNP11 is due to a preferable mRNA secondary structure. The RBS hairpin loop of pNP9 probably acts to inhibit translation while creation of another hairpin loop which exposes the initiation codon is responsible for the increased expression.

The plasmid pNP14 was constructed, such that it had the S-D to AUG sequence of pNP9, but the coding sequence of pNP11. The expression plasmid pNP14, therefore, differed from pNP9 in only three residues, those in the degenerate position of three codons, yet strains containing pNP14 expressed over ninety times more  $\mu$  than those containing pNP9. This increase in expression cannot be explained on the basis of the three residue changes optimizing codon usage, for they introduce less favoured codons, as judged by the frequency of their occurrence in strongly expressed E.coli genes.

The increased expression is best explained by the abolition of the RBS hairpin loop in pNP9. The magnitude of the increases in expression observed, and the few residue changes introduced, makes it difficult for these results to be interpreted in terms of changes in mRNA stability.

The expression of pNP8, which encodes the trpE- $\mu$  fusion protein from a native trpE RBS, was also examined, and found to express at approximately the same level as pNP14. This may indicate that the RBS of pNP14 is equivalent to that of trpE in terms of its efficiency in directing translation initiation. The native trpE RBS of pNP8 and that of pNP14 were found to have no secondary structures hindering the use of their RBS sequences.

Expression of  $\mu$  Protein in E. Coli

E. Coli B cells containing the  $\mu$  expression plasmid pNP11 were grown under inducing conditions and soluble and insoluble extracts prepared, and analysed by SDS-PAGE. A novel band was seen after staining the gel with Coomassie blue in the lane containing proteins from the insoluble fraction (see Figure 6, lane 2). This band was not seen in the negative control lane which contained proteins from the same fraction from cells harbouring pCT70 (see Figure 6, lane 3). The novel band was found to migrate to a position corresponding to a protein of a molecular weight within less than 5% of the actual molecular weight of non-glycosylated  $\mu$  of 62.5Kd. A duplicate set of lanes were transferred to nitrocellulose, and Western blotted. Alignment of the stained gel and the blot autoradiogram confirms that this novel band is antigenically related to IgM (see Figure 6, lanes 4 and 8). No band was found in extracts from cells containing pCT70 (see Figure 6, lanes 5 and 7). Only a low amount of  $\mu$  was found in the soluble fraction (see Figure 6, lane 6).

A greatly increased level of expression of  $\mu$  was found in E. Coli B compared to HB101. Pulse chase analysis demonstrated that in E. Coli B, a similar level of  $\mu$  protein was detected after a 60 minute chase (figure 7, lane 3) as was seen after the initial labelling period (Figure 7, lane 1). In HB101, however, very little  $\mu$  protein could be seen after a 10 minute chase (Figure 7, lane 7), and none after 30 minutes (Figure 7, lane 8), compared to the amount detected after the initial labelling period (Figure 7, lane 5).

For pulse chase analysis, inductions were set up as described above, except that the medium used consisted of: proline (0.3g/L), leucine (0.1g/L), Difco methionine assay medium (5g/L), glucose (60mg/L), thiamine (10mg/L),


CaCl<sub>2</sub> (22mg/L), MgSO<sub>4</sub> (0.25g/L) and carbenicillin (0.1g/L).  
During exponential growth cells were pulse labelled with 30  $\mu$ Ci/ml  
L- [<sup>35</sup>S] methionine for 2 minutes, after which unlabelled methione  
(100  $\mu$ g/ml) was added and the incubation continued for the times  
5 indicated.

Induced E. Coli B cells harbouring pNP14 when examined by phase  
contrast microscopy were found to contain inclusion bodies.

#### Expression of $\lambda$ Light Chain in E.Coli

A fresh 1 ml overnight culture of E.Coli HB101 or RV308  
containing the plasmid under study was grown in L Broth  
(Maniatis et al., Molecular Cloning, Cold Spring Harbor  
25 Laboratory, 1982), supplemented with carbenicillin to  
100  $\mu$ g/ml.

A 1:100 dilution of this culture was made into M9 medium  
(Maniatis, 1982, op.cit.) supplemented with glucose,  
vitamin B<sub>1</sub>, carbenicillin, leucine and proline and the



cultures shaken at 37° for 5-6 hours. At this time <sup>35</sup>S-methionine was added for 5-20 minutes. The cells were then harvested by centrifugation, lysed by boiling in 1% SDS for 2 minutes and diluted by addition of a buffer  
5 consisting of 2% Triton X-100, 50 mM Tris pH 8, 0.15 M NaCl and 0.1 mM EDTA. Immunoprecipitations were carried out by addition of antisera to aliquots of labelled E.Coli extracts and, after incubation at 4° overnight, immune complexes were isolated by binding to Staphylococcus  
10 aureus fixed cells. The complexes were dissociated by boiling in 60 mM Tris pH 6.8 buffer containing 10% glycerol, 3% SDS and 5% mercaptoethanol and the liberated proteins analysed on 10% or 12.5% acrylamide/SDS gels (Laemmli, UK. Nature, 227, 680-685, 1970). Gels were  
15 stained with Coomassie Brilliant blue to visualise the protein bands while labelled proteins were detected on Fuji-RX film by fluorography using 1 M sodium salicylate.

HB101 cells containing pNP3 and other plasmids were grown under inducing conditions to an OD<sub>600</sub> = 0.6 and the proteins present examined by specific immunoprecipitation  
20 substantially as described above. The results obtained are given in Figure 8, in which lanes 1 and 2 respectively were extracts from pCT54-19 and pNP3, immunoprecipitated with normal rabbit serum; lanes 3, 4 and 5 were extracts immunoprecipitated with rabbit anti-lambda serum and  
25 represent pNP4 (lane 3), pNP3 (lane 4) and pCT54-19 (lane 5). The position of unlabelled lambda protein from MOPC104E is indicated on the left hand side of Figure 8. All of the plasmids produced a protein which reacted with  
30 rabbit anti-mouse  $\lambda$  light chain serum and comigrated with authentic  $\lambda_1$  light chain from the mouse myeloma MOPC104E. pNP3 however produced the most of this protein (compare lane 4 with lanes 3 and 5 in Figure 8). No such band was detected with control immunoprecipitations using normal  
35 rabbit serum (Figure 8, lanes 1 and 2). Studies using brief pulses with <sup>35</sup>S-methionine followed by chasing with

excess cold methionine indicated that the recombinant  $\lambda$  light chain had a 1/2 life of about 20 minutes.

In vitro transcription/translation (Pratt et al, Nucleic Acids Research, 9, 4459-447, 1981) confirmed that pNP3  
5 coded for a protein which comigrated with authentic  $\lambda$  light chain. This 25 Kd product was synthesised in vitro at a rate comparable to that of  $\beta$ -lactamase indicating that it is synthesised in vivo at a level of 0.5% of total E.Coli proteins. This figure is in good agreement with  
10 the percentage of recombinant  $\lambda$  product synthesised in vivo (0.4-2%) as determined from the following equation:

$$\% \text{ specific CPM} = \frac{100 \times (\text{CPM with anti-}\lambda) - (\text{CPM with normal serum})}{\text{total CPM}}$$

The procedure used to disrupt cells was as follows (Emtage  
15 J S et al, Proc. Natl. Acad. Sci., 80, 3671-3715, 1983). E.Coli HB101/pNP3 grown under inducing conditions were harvested and resuspended in 0.05 M Tris pH 8, 0.233 M NaCl, 5% glycerol containing 130 g/ml of lysozyme and incubated at 4°C or room temperature for 20 minutes. Sodium  
20 deoxycholate was then added to a final concentration of 0.05% and 10  $\mu$ g of DNAase 1 (from bovine pancreas) was added per g wet wt of E.Coli. The solution was incubated at 15°C for 30 minutes by which time the viscosity of the solution had decreased markedly. The extract was centri-  
25 fugal (at 10,000 x g for 15 minutes for small volumes (1 ml) or 1 hour for larger volumes) to produce a soluble and an insoluble fraction.

For immunoprecipitations, the soluble fraction was diluted  
30 in the Triton-containing buffer described above and the insoluble fraction solubilised by boiling in 1% SDS followed by dilution in Triton-containing buffer. HB101 cells containing pNP3 or pNP4 were grown under inducing

- 30 -

conditions, pulse labelled with [ $^{35}$ S] methionine, separated into soluble and insoluble fractions and analysed by sodium dodecyl sulfate polyacrylamide gel electrophoresis. The results obtained are shown in Figure 9 in which lanes 1 and 3 correspond to the soluble fraction from HB101-pNP3 immunoprecipitated with normal rabbit serum and rabbit anti-lambda serum respectively; lanes 2 and 4 correspond to the insoluble fraction from HB101-pNP3 immunoprecipitated with normal rabbit serum and rabbit anti-lambda serum respectively, lanes 5 and 7 correspond to the soluble fraction from HB101-pNP4 immunoprecipitated with normal rabbit serum and rabbit anti-lambda serum respectively; lane 9 corresponds to the soluble fraction from HB101-pNP3, and lane 10 corresponds to the insoluble fraction from HB101-pNP3. The position of unlabelled lambda protein from MOPC104E is indicated on the left hand side of Figure 9. When such a procedure was carried out using pNP3, the recombinant  $\lambda$  light chain protein was present in the insoluble fraction (Figure 9, lane 4) rather than the soluble fraction (Figure 9, lane 3). No 25 Kd band comigrating with authentic  $\lambda_1$  light chain was present when either the soluble or insoluble fraction were immunoprecipitated using normal rabbit serum (Figure 9, lanes 1 and 2). By inference it is likely that all Ig light chains will fractionate as in this specific example.

In the absence of specific immunoprecipitation a novel protein band was not visible from extracts of HB101 containing pNP3 nor was the  $\lambda$  protein found to accumulate. However, there was a dramatic difference  
5 when pNP3 was induced in the K12 strain E103s. In this strain  $\lambda$  protein was found to accumulate during induction until the cells reaches stationary phase (figure 10, lanes 3-5) to a level of about 150 times that found in HB101 as determined by an ELISA (enzyme  
10 linked immunosorbent assay). These cells were found to contain inclusion bodies which appeared refractile under light microscopy, a phenomenon characteristic of high level expression of foreign proteins. An estimate of the percentage of total E. Coli protein  
15 represented by recombinant  $\lambda$  protein was obtained by separating the proteins by gel electrophoresis, staining them, with Coomassie blue and scanning the stained gel with a Joyce-Loebl chromoscan 3. This  
20 method showed that  $\lambda$  was the major protein present (Figure 10, lane 5) and represented 13% of total E. Coli protein. The  $\lambda$  protein had a half-life of 20 minute in HB101 but accumulated to very high levels in E103s, suggesting that Lambda protein was much more  
25 stable in the latter strain. After cell lysis and centrifugation of HB101 or E103s containing pNP3,  $\lambda$  light chain was detected in the insoluble (Figure 10, lanes 7 and 10) but not in the soluble fractions (Figure 10, lanes 6), as determined by Coomassie blue  
30 staining. The identity of the major Coomassie blue stained band as  $\lambda$  protein was confirmed by Western blot analysis (Figure 10, lanes 8-10). The presence of such immunoreactive bands was specific to pNP3 containing cells. When extracts from cells containing pCT70, a prochymosin expressing plasmid, were subjected  
35 to the same analysis no bands were detected. This more sensitive technique showed that a small amount of the  $\lambda$  protein was in the soluble fraction (Figure 10, lane 9). The presence of a number of distinct immuno-



reactive proteins all smaller than full-length protein were also detected. These may result from proteolytic degradation of  $\lambda$  protein, from premature termination of transcription or from internal initiation of translation.

Expression of  $\mu$  and Lambda Polypeptides in the same Bacterial Cell

Each of the Ig  $\mu$  and  $\lambda$  genes in expression plasmids were transformed into the same E.Coli cell to direct the synthesis of both Ig  $\mu$  and  $\lambda$  polypeptides. In order to overcome plasmid incompatibility and provide a second antibiotic resistance marker, the trp promoter and  $\lambda$  sequences were excised from pNP3 on a Hind III - Bam HI fragment and inserted into the Hind III - Bam HI fragment of pACYC 184. The resultant plasmid pACYC  $\lambda$  caused the E.Coli to grow very poorly. This weak growth was thought to result from read-through of RNA polymerase into the origin of replication. However, the inhibition of growth was virtually eliminated by the cloning in of the bacteriophage T7 transcriptional terminator at the Hind III site of plasmid pACYC  $\lambda$ .

This terminator functions in both orientations. The resultant plasmid pAC  $\lambda$  T7-1 has a chloramphenicol resistance gene and an origin compatible with the pBR 322-derived origin on pNP14, the Ig  $\mu$  expressing plasmid. Transformation of both plasmids in the same E.Coli B was achieved in two steps. Firstly pNP14 was introduced followed by pAC  $\lambda$  T7 - I in two sequential transformations to give ampicillin and chloramphenicol resistant clones.

E.Coli B cells derived from this double-transformant clone showed the presence of inclusion bodies and two novel polypeptide bands on stained gels of the insoluble fraction after lysis. These two bands correspond both with immunological activity by Western blotting for Ig  $\mu$  and Ig  $\lambda$  and in their expected molecular weights. It has thus been shown that the double-transformant clone expresses both the heterologous genes. This had not hitherto been shown.

The presence of  $\lambda$  light chain in the insoluble fraction was a useful purification step since it both concentrated the protein and separated it from the bulk of E. Coli soluble proteins.

5

For further purification of the  $\lambda$  light chain, the cell debris were dissolved in 10mM Tris-HCl pH8.0, 25% formamide, 7M urea, 1mM EDTA and 2mM dithiothreitol. This material was loaded onto a DEAE Sephacel column (Pharmacia) (1x 25cm at a flow rate of 5ml/hr) which had been equilibrated in 9M urea, 10mM Tris-HCl pH8.0, 1mM EDTA and 2mM DTT. The DEAE Sephacel column was developed using a 0-150mM NaCl gradient in loading buffer. The eluted peak  $\lambda$  light chain immunoreactivity, corresponding to the major peak of protein, was diluted to a final concentration of 2.25M urea, 10mM Tris-HCl pH8.0, 1mM EDTA, 2mM DTT and loaded onto an octyl-Sephacel column (Pharmacia) (2.5 x 10cm). Material was eluted by use of a urea gradient of 2.25-9M urea. The peak material was pooled, dialysed into ammonium bicarbonate and lyophilised. Following this step, only a single band of Coomassie blue stainable material corresponding to recombinant  $\lambda$  protein was visualised by SDS-PAGE.

25

The  $\mu$  heavy chain was purified from 9M urea solubilised pellets by anion exchange chromatography and chromatofocussing (Pharmacia).

30

It was of great interest to determine whether the concomitant expression of  $\mu$  and  $\lambda$  would lead to the formation of functional IgM. In order to determine this, extracts were made from E. Coli containing both Ig  $\mu$  and  $\lambda$  polypeptides and these tested for antigen binding. We used a two-site sandwich ELISA which detects  $\mu$  chain binding to haptenated bovine serum albumin (NIP-caproate-BSA).

In the NIP binding assay, bovine serum albumin (BSA) is reacted with an equimolar amount of NIP-Cap-N-hydroxy-succinamide at pH 7.5 in 10mM phosphate buffer. The resulting NIP-Cap-BSA is separated from free  
5 NIP-Cap on a G-50 Sephadex column.

Microtitre plates (96 well Nunc Immuno Plate 1) are coated with 100  $\mu$ l of a solution of 10  $\mu$ g/ml NIP-BSA in sodium carbonate/sodium bicarbonate 0.1M, pH 9.6  
10 buffer (coating buffer) overnight at 4°C.

The coated plates are then blocked for non-specific binding by addition of 100  $\mu$ l of 0.5% casein in coating buffer (blocking buffer) and incubating at 37°C for  
15 1 hour.

The coated and blocked plates are then washed three times in 150 mM NaCl, 0.05% NP40, 20mM Tris-HCl, pH 7.8-buffer (washing buffer).  
20

The washed plates are shaken free of excess washing buffer and samples are added in their buffer or typically in 0.5% casein in washing buffer (sample buffer) to a 100  $\mu$ l final volume.  
25

In order to demonstrate the nature of the binding to NIP-BSA, samples for testing were supplemented with either NIP or NP in free solution at various concentrations (from 60 to 0.3  $\mu$ M for NIP or from 600 to 3  $\mu$ M  
30 for NP).

The plates with samples are then incubated at 37°C for one hour and thereafter washed three times with washing buffer.  
35

The washed plates are then inoculated with 100  $\mu$ l of anti  $\mu$ -peroxidase conjugate (1:1000 dilution; TAGO Inc) in sample buffer, and incubated for one hour at 37°C. The plates are then washed three times in washing  
5 buffer.

The washed plates are inoculated with 100  $\mu$ l of 0.1M Na acetate-citrate, pH 6.0, 0.1 mg/ml tetramethylbenzidine, 13mM  $H_2O_2$  (peroxidase substrate) and  
10 incubated at room temperature for one hour. The reaction is terminated by the addition of 25 $\mu$ l of 2.5M  $H_2SO_4$ .

The microtitre plates are then read in a plate reader  
15 (Dynatech) at 450 nm with a reference of 630 nm. The  $A_{450}$  was related to the level of a standard protein, B1-8 anti-NIP IgM.

This assay demonstrates sensitivity to 60pg of B1-8 IgM.  
20 The extracts were prepared as soluble and insoluble material. The insoluble material was solubilized in the same buffer used in lysis but containing 8M urea followed by its dilution for assay.

25 In order to obtain activity for the Ig  $\mu$  and  $\lambda$ , extracts were made of the insoluble fraction and these dialysed into buffer conditions in which disulphide interchange will occur at a higher frequency.

30 Production of functional antibodies from E. Coli expressing both heavy and light chains was achieved by lysing the cells and clarifying the supernatant by centrifugation. The insoluble material was washed, followed by sonication (3 times for 3 minutes), and  
35 finally dissolved in 9M urea, 50mM glycine-NaOH pH10.8,

1mM EDTA, and 20mM 2-mercaptoethanol. This extract was dialysed for 40 hours against 3 changes of 20 vols. of 100mM KCl, 50mM glycine-NaOH pH10.8, 5% glycerol, 0.05mM EDTA, 0.5mM reduced glutathione and 0.1mM oxidised glutathione. The dialysate was cleared by centrifugation at 30,000g for 15 minutes and loaded directly onto DEAE Sephacel, followed by development with a 0-0.5M KCl linear gradient in 10mM Tris-HCl, 0.5mM EDTA, pH8.0.

10 The purified Ig $\mu$  and  $\lambda$  were treated as above, except that no anion exchange chromatography was carried out. The preparation was finally dialysed into phosphate buffered saline, 5% glycerol, 0.01% sodium azide and 0.5mM EDTA pH7.4.

15 The results from assays of material processed in this way indicated that some activity was obtained. The level of activity obtained in this way was too low to do any detailed studies on, so the resultant dialysate was purified by anion exchange chromatography (Figure 11). This process resulted in the isolation of significant NIP-cap-BSA binding activity over that of background binding to BSA (Figure 11). The assay of the fractions for the level of Ig $\mu$ , expressed as B1-8 IgM equivalents demonstrated two peaks of activity. This was not found to correlate with full length Ig $\mu$  by Western blotting. The first peak observed may represent a fragment of Ig $\mu$ . The separation of NIP-CAP-BSA binding activity from the majority of full length Ig $\mu$  and protein indicates that the hapten binding activity is contained within a particular molecular species formed at low efficiency.

The processing of insoluble material obtained from Ig  
35  $\mu$  expression in E. Coli produced a similar IgM protein

profile but without NIP-cap-BSA binding activity.  
This demonstrates that the activity recovered was a  
property of the combined immunoglobulin expression,  
not of some E. Coli factor, or of the Ig  $\mu$  heavy chain  
5 alone.

Further studies of the characteristics of the hapten  
(NIP-cap-BSA) binding were carried out. As samples  
were diluted they showed less binding to hapten in a  
10 very similar way to the original antibody (Figure 12).  
Free hapten was found to inhibit most of the binding  
activity in both undiluted and diluted samples. Using  
B1-8 antibody as a standard for both IgM and hapten  
binding, the specific activity of the assembled antibody  
15 was calculated to be  $1.4 \times 10^4$  gm/gm of IgM equivalents.  
This value demonstrates the inefficient recovery of  
activity, but possibly represents an underestimate  
of the specific activity due to an overestimate of full-  
length Ig  $\mu$  in these fractions, as described above.

20

#### Heteroclitic Nature of Recombined Antibody

Detailed specificity of binding to NIP-cap-BSA was  
investigated by comparing the assembled antibodies  
with B1-8 IgM in the presence of free NIP-cap and NP-  
25 cap (Figure 13). Both B1-8 IgM and the assembled anti-  
bodies showed that higher NP-cap than NIP-cap concentra-  
tions were required to inhibit NIP-cap-BSA binding.

The heteroclitic nature is demonstrated by the molar  
30 ratio of NIP to NP at 50% inhibition. The concentrations  
of NIP and NP at 50% inhibition (I50) were found to  
be similar for both B1-8 and the assembled antibodies  
as shown in Table 2. Also the NP I50/NIP I50 ratios  
were similar (Table 2).

35

Table 2. Hapten concentration at 50% inhibition (I50) of binding of antibodies to NIP-cap-BSA solid phase.

SD = Standard Deviation.

5

	<u>NIP I50 <math>\mu</math>M</u>	<u>NP I50 <math>\mu</math>M</u>	<u>NP I50</u> <u>NIP I50</u>
B1-8 IgM	0.13 (SD, 0.05)	3.7 (SD, 2.9)	29
Fraction 26	0.34 (SD, 0.09)	1.9 (SD, 0.4)	6
10 Fraction 27 and 28	0.11 (SD, 0.02)	1.1 (SD, 0.3)	10
Purified $\mu$ and $\lambda$	0.4	0.84	22

It has thus been shown that not only is it possible to express  $\mu$  and  $\lambda$  proteins having intact variable domains, but also that it is possible to express both  $\mu$  and  $\lambda$  proteins on compatible plasmids in the same cell. This latter has not hitherto been disclosed or even suggested. Moreover, it proved possible to derive functional Ig molecules from E. Coli cells which expressed both  $\mu$  and  $\lambda$  proteins.

It has thus been shown that the process of the present invention produces a functional Ig molecule by recombinant DNA techniques.

25

Construction of Expression Plasmids for Yeast

The plasmids used for expression in yeast (Saccharomyces cerevisiae) are based on pBR322 and the yeast 2 micron plasmid. These plasmids are the subject of copending European Patent Application EP 0 073 635, the disclosure of which is incorporated hereby reference. The yeast phosphoglycerate kinase gene (PGK) provides the 5' sequence necessary for initiation of transcription and insertion of genes can be made at the unique BglII site in the plasmid pMA 3013 described in the above mentioned European patent application. This plasmid has now been renamed pMA91 as is so referred to hereinafter.

a) met-mu ( $\mu$ )

BglII and BclI produce compatible 5'-GATC ends. The met- $\mu$  gene from pNP2 was excised on a partial BclI fragment and ligated into the unique BglII site of pMA91.

b) Pre-mu ( $\mu$ )

The plasmid pCT54 u containing the full length pre- $\mu$  cDNA was digested with Hind III. This cuts at the 3' side of the  $\mu$  cDNA. This Hind III site was changed to a BclI site by incubation with  $T_4$  DNA polymerase in the presence



of all 4 nucleotides. The linker R107 of sequence  
TTTTGATCAAAA, which contains an internal BclI was ligated  
to the DNA obtained above. After ligation one aliquot  
of the resultant DNA was digested with BclI and AccI and  
5 the  $\mu$  AccI-BclI fragment was isolated from the resultant  
mixture by gel electrophoresis. A separate aliquot of  
the resultant DNA was digested with MboII and ligated  
with chemically synthesised oligonucleotide linkers R121  
and R112 which have cohesive BclI and MboII ends.

10 R121 5' GATCAATGGGATGGAGCTGT 3'  
R112 5' TTACCCTACCTCGAC 3'

The ligation reaction was terminated and the resultant DNA  
digested with AccI to give a  $\mu$  MboII-AccI fragment which  
was isolated from the resultant mixture by gel electro-  
phoresis on a 5% polyacrylamide gel.

15 The two  $\mu$  fragments obtained above were ligated together  
and then digested with BclI to eliminate unwanted ligation  
products after which the resultant pre- $\mu$  DNA was ligated to  
BglIII cut plasmid pMA91.

The resultant ligation mix was used to transform E.Coli  
20 strain HB101 to ampicillin resistance. A colony of the  
transformed bacteria was isolated and was found to contain  
a plasmid exhibiting the predicted enzyme digestion pattern.  
The 5' end of the inserted DNA from this plasmid was se-  
quenced and shown to have the anticipated sequence for a  
25 pre- $\mu$  cDNA.

c) met-lambda ( $\lambda$ )

The met- $\lambda$  DNA was cloned into the BglIII site of plasmid pMA91.  
Plasmid pCT54 Clone 1 was cut with Hind III to the 3' side  
of the met- $\lambda$  cDNA and this cleavage site was altered to

a BclI site in a similar manner as previously described for pNP2. Following this the plasmid DNA was cut 5' of the  $\lambda$  gene at the BclI site and the resultant  $\lambda$  DNA was cloned into the BglII site of pMA91.

5 d) Pre-lambda ( $\lambda$ )

Pre- $\lambda$  cDNA was reconstructed into pMA.91 as follows. The plasmid pCT54 was digested with BclI and Hind III and the resultant vector DNA was isolated by gel electrophoresis and ligated together with two synthetic oligonucleotides,  
10 R162 and R163, and the two fragments from plasmid pAT  $\lambda$  1-15.

R162 5' GATCAATGGCCTGGATT 3'

R163 5' GTGAAATCCAGGCCATT 3'

These fragments were produced by digesting pAT  $\lambda$  1-15 with FokI and Hind III and isolating the 5' 300 base pair FokI fragment and the 3' 600 base pair FokI-Hind III fragment  
15 by gel electrophoresis. The resultant ligation mixture was used to transform E.Coli strain HB101 to ampicillin resistance and a colony of bacteria containing a plasmid having the predicted correct restriction enzyme digestion pattern was isolated. The 5' end of the inserted DNA was  
20 sequenced and found to have the anticipated sequence of a pre- $\lambda$  cDNA. This plasmid was digested with Hind III and the resultant Hind III cleaved ends converted to a BclI site by blunting with  $T_4$  polymerase and ligating with the linker R107. The resultant plasmid was then digested  
25 with BclI and the pre- $\lambda$  cDNA isolated on a BclI fragment by gel electrophoresis. This BclI fragment was then ligated to BglI cut pMA91 to give a pMA 91 plasmid containing the full length pre- $\lambda$  cDNA.

Expression of Immunoglobulin Genes in Yeast

30 The pMA91 derivative plasmids containing the pre- $\lambda$  and pre- $\mu$  genes as prepared above were used to transform

Saccharomyces cerevisiae yeast host organisms and the pre- $\lambda$  and pre- $\mu$  genes were expressed by the transformed cells.

Saccharomyces cerevisiae strain MD46 when transformed  
5 with pMA91 pre- $\lambda$  or pMA91 pre- $\mu$  gave rise to colonies  
which expressed immunoreactive proteins as revealed on  
Western blots. Yeast cells containing plasmid pMA91  
pre- $\lambda$  produced a protein which reacted with anti- $\lambda$   
antiserum and co-migrated on polyacrylamide gels with  
10 bacterially synthesised mature  $\lambda$ . Similarly yeast  
cells containing pMA 91 pre- $\mu$  produced a protein which  
reacted with anti- $\mu$  antiserum and co-migrated on poly-  
acrylamide gels with bacterially synthesised mature  $\mu$   
protein. These observations indicate that both pre- $\lambda$   
15 and pre- $\mu$  are being processed to the corresponding mature  
proteins within the yeast host cell environment.

Additionally the  $\mu$  protein product (but not the  $\lambda$  protein)  
was shown to be glycosylated. Cells were incubated in  
the presence or absence of tunicamycin at a concentration  
20 of 15  $\mu\text{g/ml}$ . This compound specifically arrests the  
N-linked glycosylation of proteins. Cell extracts derived  
from cells incubated in the absence of tunicamycin showed  
higher molecular weight bands (as revealed by Western  
blotting) whilst extracts from cells incubated in the  
25 presence of tunicamycin showed no such higher molecular  
weight bands.

When cells were lysed with glass beads and the soluble and  
insoluble fractions examined it was found that the  $\mu$  and  
 $\lambda$  proteins were exclusively in the insoluble fractions,  
30 as determined by Western blotting.

### Secretion

After incubation transformed yeast cells were spun down and 1 ml volumes of supernatant were removed and passed through BSA-coated 0.2  $\mu$ M Millipore filters to remove  
5 any remaining cellular material. ELISA assays for  $\mu$  and  $\lambda$  protein were then carried out on the filtered supernatants. Only supernatants from pMA91 pre- $\lambda$  harbouring cells showed detectable levels of immunoglobulin protein. Increased amounts of  $\lambda$  protein was  
10 detected in the supernatant when cells were grown in minimal medium to OD<sub>660</sub> = 0.2 and then spun down and resuspended in YPAB, and harvested at OD<sub>660</sub> = 1.0-1.5. Thus the  $\lambda$  protein but not the  $\mu$  protein is secreted from yeast cells.

### 15 Intracellular Location

After incubation cells containing pMA91 pre- $\mu$  were converted to spheroplasts and fixed in 5% acetic acid/95% ethanol (v/v) and incubated with fluorescein conjugated goat anti-mouse  $\mu$ . The fixed cells were examined by  
20 fluorescent microscopy and the  $\mu$  protein was found to be localised in the periphery of the spheroplasts and especially in vacuoles.

### Expression of both $\mu$ and $\lambda$ Proteins in the Same Cells

In order to express both genes in the same host cells it  
25 was necessary to provide compatible plasmids for use in transformation. The pre- $\lambda$  gene was excised from pMA91 pre- $\lambda$  on a Hind III fragment and inserted into the Hind III site of plasmid pLG89 (Griti L, and Davies J, Gene, 25 179-188, 1983). This plasmid contains a ura3 marker  
30 which can be used as a positive selection for transformed host cells. A convenient ura3<sup>-</sup> host organism is S. cerevisiae strain X4003-5B. Both plasmids, pMA91 pre- $\mu$

and pLG89 pre- $\lambda$  , were transformed into this strain, and colonies were grown which contained together both the ura3 and leu2 markers. After incubation of the transformed cells both  $\lambda$  and  $\mu$  proteins together were  
5 detected in the same cultures of X4003-5B using ELISA techniques. Levels of expression were comparable to those obtained for the individual genes in MD46 cells and for the  $\lambda$  gene alone in X4003-5B.

In addition, to check for assembly of the  $\lambda$  and  $\mu$  proteins in vivo the following procedure was followed.  
10

After the growth to an  $OD_{660} \cong 1$ , the transformed X4003-5B cells were spun down and resuspended in buffer, either 5 mM borate buffer at pH 8.0 or phosphate buffer at pH 5.8 either with or without detergent (e.g. 0.5% Triton X).  
15 The suspended cells were then lysed by vortexing with glass beads in buffer as above and the insoluble material was spun down. After centrifugation aliquots of the supernatant were assayed in a NIP binding assay (as described previously for assembled E.Coli  $\mu$  and  $\lambda$  proteins).  
20 Specific antigen binding activity was detected in the supernatant and this activity was shown to be specifically competed out by free NIP (the specific antigen). These results indicated that  $\mu$  and  $\lambda$  protein were expressed within the X4003-5B cells and assembled into  
25 functional immunoglobulin molecules in vivo.

In the above experiments the culture medium used for incubating the transformed cells was yeast minimal medium (containing, per litre, 6.7 g DIFCO yeast nitrogen base without amino acids, 10 g glucose, and 200 mg each of  
30 histidine, tryptophan, methionine and adenine). When the  $\lambda$  protein alone was being expressed the medium contained in addition 200 mg/l of leucine, and when the  $\mu$  protein alone was being expressed the medium contained in addition 200 mg/l of uracil.

## CLAIMS:

1. A process for producing a heterologous multi chain polypeptide or protein in a single host cell, which comprises transforming the host cell with DNA sequences coding for each of the polypeptide chains and expressing  
5 said polypeptide chains in said transformed host cells.
2. A process according to claim 1, wherein each DNA sequence is inserted individually into a vector and the individual vectors are used to transform the single host cell.
- 10 3. A process according to claim 1, wherein at least two of the DNA sequences are inserted into separate parts of a vector which is used to transform the single host cell.
4. A process according to claim 2 or claim 3 wherein  
15 the or each vector is a plasmid.
5. A process according to claim 4, wherein the plasmid is col E1, pcR1, pBR322, RP4 or phage DNA.
6. A process according to any one of claims 1 to 5 wherein the host cell is a bacterium or a yeast.
- 20 7. A process according to claim 6, wherein the host cell is E.Coli, B. subtilis or S.cerevisiae.
8. A process according to claim 7, wherein the host cell is E.Coli strain HB101, X1776, X2887, PS410, MRC 1 RV308, 2 or E1035.

9. A process according to any one of claims 1 to 8, wherein the polypeptide chains are expressed and secreted by the host cell.

5 10. A process according to any one of claims 1 to 8, wherein the polypeptide chains expressed by the host cell are solubilised under conditions which facilitate assembly of the active multi chain polypeptide or protein.

10

11. A process according to any one of claims 1 to 10, for producing an Ig molecule or an immunologically functional Ig fragment, wherein the host cell is transformed with DNA sequences coding for at least  
15 the variable domains of the Ig heavy and light chains.

12. A process according to claim 11, wherein the DNA sequences code for the complete Ig heavy and light chains.

20

13. A process according to claim 11 or claim 12, for producing an Ig molecule or fragment having at least one constant domain, wherein the constant domain is derived from the same source as the variable domain  
25 to which it is attached.

14. A process according to claim 11 or claim 12, for producing an Ig molecule or fragment having at least one constant domain, wherein the constant domain  
30 is derived from a different source for that from which the variable domain to which it is attached is derived.

15. A process according to any one of claims 11 to 14,  
35 wherein the DNA coding sequences for the Ig molecule

or fragment are derived from one or more monoclonal antibody producing hybridomas.

- 5 16. A process for increasing the level of protein expression in a transformed host cell comprising altering the vector DNA sequence in the region of the initiator ATG codon to one which substantially eliminates RNA secondary structure.
- 10 17. A process according to claim 16 wherein the DNA sequence is altered to minimise the absolute value of the  $\Delta G$  value of the sequence.
- 15 18. A heterologous multi chain polypeptide or protein produced by recombinant DNA technology from a single host cell.
- 20 19. An Ig molecule or an immunologically functional Ig fragment produced by recombinant DNA technology from a single host cell.
- 25 20. An Ig molecule or fragment according to claim 19, including at least one constant domain derived from the same source as the variable domain to which it is attached.



21. An Ig molecule or fragment according to claim 19, including at least one constant domain derived from a source different from the source from which the variable domain to which it is attached was derived.
- 5 22. An Ig fragment according to claim 20 or claim 21, comprising only one constant domain attached to each variable domain.
23. An Ig molecule according to claim 20 or claim 21, comprising sufficient constant domains attached to each  
10 variable domain to produce a complete Ig molecule.
24. An Ig heavy or light chain or a fragment thereof having a complete variable domain, as a product of recombinant DNA technology.
25. A vector for use in a process according to any one  
15 of claims 1 to 17.
26. A vector for use in transforming a host cell to enable it to produce a molecule according to any one of claims 18 to 24.
27. A vector according to claim 25 or claim 26, which is  
20 a plasmid
28. A host cell transformed with a vector according to any one of claims 25 to 27.
29. A therapeutic composition comprising a molecule according to any one of claims 18 to 24 and a pharmaceuti-  
25 cally acceptable carrier.
30. A process for preparing a composition according to claim 29, comprising mixing a molecule according to any one of claims 18 to 24 with a pharmaceutically acceptable carrier.
- 30 31. An affinity chromatography medium comprising a molecule according to any one of claims 18 to 24 immobilised on a chromatographic medium.
32. An immunological assay method in which the or each antigen binding Ig molecule or fragment is a molecule or  
35 fragment according to any one of claims 19 to 24.

0120694

33. An Ig molecule or fragment according to any one of claims 18 to 24 having attached thereto a cytotoxic, enzymic or structural polypeptide moiety synthesised in the cell with the Ig molecule or fragment





FIG. 1

2/13



Legend.

E = E corRI

H = HinfI

H3= Hind III

FIG. 2

3/13




FIG. 3

4/13

Possible 2 structures of  $\mu$  mRNAs

5'→3'







FIG. 4

5/13




FIG. 5

6/13




FIG. 6



7/13




FIG. 7




FIG. 8

9/13




FIG. 9




FIG. 10

11/13




FIG. 11

12/13





FIG. 12



Binding of antibodies to NIP-cap BSA  
B1-8 IgM (■), fraction 26 (▲),  
purified Ig  $\mu$  or  $\lambda$  (●), in the presence  
of free NIP-cap(---) or NIP-cap (—).

FIG. 13